arXiv:hep-ph/9910240v1 5 Oct 1999

Resummation for direct photon and W + jet production

Nikolaos Kidonakis

Department of Physics, Florida State University,
Tallahassee, FL 32306-4350, USA

E-mail: kidonaki@sg1.hep.fsu.edu

Abstract

I discuss the resummation of soft gluon contributions to direct photon and W + jet production and I present some results for the next-to-next-to-leading order expansions of the resummed cross sections for these processes near threshold.

1. Introduction

Direct photon production and W + jet production are processes of great interest for a number of reasons. Direct photon production is important for determinations of gluon distributions while W + jet production is relevant for estimating backgrounds to new physics. Near threshold for the production of the final state in these processes one can resum logarithmic corrections originating from soft gluon emission to all orders in perturbative QCD. These corrections are potentially large. A formalism for the resummation of soft-gluon contributions at next-to-leading logarithmic (NLL) accuracy has been recently developed for a variety of QCD hard scattering processes, including heavy quark, jet, direct photon, and W + jet production (for reviews see Refs. [1, 2]). The resummed cross sections have been expanded up to next-to-next-to-leading order (NNLO), thus providing new analytical higher order predictions, and exhibit a substantially reduced scale dependence relative to next-to-leading order (NLO); this is expected on theoretical grounds [3].

2. Direct photon production

First, we discuss resummation for direct photon production [4, 5, 1, 6, 7, 8] in hadronic processes

$$h_1 + h_2 \to \gamma + X\,. \tag{1}$$

The factorized cross section may be written as a convolution of parton distributions ϕ with the partonic hard scattering $\hat{\sigma}$

$$\sigma_{h_1 h_2 \to \gamma} = \sum_f \phi_{f_1/h_1} \otimes \phi_{f_2/h_2} \otimes \hat{\sigma}_{f_1 f_2 \to \gamma}\,. \tag{2}$$

Near partonic threshold $\hat{\sigma}$ includes large logarithmic terms that can be resummed to all orders in perturbation theory. If we take moments of the above equation, with N the moment variable, we can write the partonic cross section as

$$\begin{aligned}\tilde{\sigma}_{f_1 f_2 \to \gamma}(N) &= \tilde{\phi}_{f_1/f_1}(N)\, \tilde{\phi}_{f_2/f_2}(N)\, \hat{\sigma}_{f_1 f_2 \to \gamma}(N) \\ &= \tilde{\psi}_{f_1/f_1}(N)\, \tilde{\psi}_{f_2/f_2}(N)\, \tilde{J}(N)\, H\, \tilde{S}(N)\,, \end{aligned} \tag{3}$$

where in the second line of Eq. (3) we have introduced a refactorization in terms of new center-of-mass parton distributions ψ and a jet function J that absorb the collinear singularities from the incoming partons and outgoing jet [9, 10, 11, 4], respectively; a soft gluon function S that describes noncollinear soft gluon emission [10, 11]; and a short-distance hard scattering function H.

We may then solve for $\hat{\sigma}_{f_1 f_2 \to \gamma}(N)$ in Eq. (3). After resumming the N-dependence of all the functions in the above equation, we may write the resummed cross section as

$$\begin{aligned}\hat{\sigma}_{f_1 f_2 \to \gamma}(N) &= \exp\left\{ \sum_{i=1,2} \left[E^{(f_i)}(N_i) + E^{(f_i)}_{\rm scale} \right] \right\} \\ &\times \exp\left\{ E'_{(f_J)}(N) \right\}\, H\left(\alpha_s(\mu^2)\right)\, S\left(1, \alpha_s(S/N^2)\right) \\ &\times \exp\left[\int_\mu^{\sqrt{S}/N} \frac{d\mu'}{\mu'}\, 2\,{\rm Re}\Gamma_S\left(\alpha_s(\mu'^2)\right) \right]\,. \end{aligned} \tag{4}$$

The exponent $E^{(f_i)}$ resums the ratio ψ/ϕ while $E'_{(f_J)}$ resums the N-dependence of the outgoing jet [11, 4]. The dependence on the scale is given by $E^{(f_i)}_{\rm scale}$. Finally, ${\rm Re}\Gamma_S$ is the real part of the soft anomalous dimension Γ_S which is determined from renormalization group analysis of the soft function S [10, 11] and has been calculated explicitly at one-loop [4, 1]. Because of the simpler color structure, Γ_S is here a simple function, in contrast to heavy

quark and jet production where it is a matrix in color space [10, 11, 12].

Since the direct photon production cross section has been calculated only up to NLO, expansions of the resummed result to higher orders provide new analytical predictions. Here we present the NNLO expansion for the partonic subprocess

$$q(p_1) + g(p_2) \to \gamma(p_\gamma) + q(p_J)\,. \tag{5}$$

We define the kinematical variables $s = (p_1 + p_2)^2$, $t = (p_1 - p_\gamma)^2$, $u = (p_2 - p_\gamma)^2$, and $v \equiv 1 + t/s$. The threshold region is given in terms of the variable $w \equiv -u/(s+t)$ by $w = 1$. Then at nth order in α_s we encounter plus distributions of the form $[\ln^m(1-w)/(1-w)]_+$ with $m \le 2n-1$.

The NLO soft gluon corrections in the $\overline{\rm MS}$ scheme for this partonic channel are given by [4, 8]

$$\begin{aligned} vw(1-v)s\frac{d\hat{\sigma}^{\rm NLO}_{qg\to q\gamma}}{dv\,dw} &= \sigma^B_{qg\to q\gamma}\frac{\alpha_s(\mu_R^2)}{\pi} \\ &\times \left\{ (C_F + 2C_A)\left[\frac{\ln(1-w)}{1-w}\right]_+ \right. \\ &+ \left[C_F\left(-\frac{3}{4} + \ln v\right) - C_A \ln\left(\frac{1-v}{v}\right)\right. \\ &\left.\left. -(C_F + C_A)\ln\left(\frac{\mu_F^2}{s}\right)\right]\left[\frac{1}{1-w}\right]_+ \right. \\ &\left. + \mathcal{O}(\delta(1-w))\right\}\,, \end{aligned} \tag{6}$$

with μ_F and μ_R the factorization and renormalization scales. Here the Born term is [13]

$$\sigma^B_{qg\to q\gamma} = \frac{1}{N_c}\pi\alpha\alpha_s(\mu_R^2)e_q^2 T_{qg} v\,, \tag{7}$$

where $T_{qg} = 1 + (1-v)^2$. Agreement is found with the exact NLO cross section in Ref. [13]. All $\delta(1-w)$ terms can be obtained by matching to the exact NLO result.

The NNLO $\overline{\rm MS}$ soft gluon corrections for $qg \to q\gamma$ are [7, 8]

$$\begin{aligned} vw(1-v)s\frac{d\hat{\sigma}^{\rm NNLO}_{qg\to q\gamma}}{dv\,dw} &= \sigma^B_{qg\to q\gamma}\frac{\alpha_s^2(\mu_R^2)}{\pi^2} \\ &\times \left\{ \frac{1}{2}(C_F + 2C_A)^2\left[\frac{\ln^3(1-w)}{1-w}\right]_+ \right. \\ &+ \left[\frac{3}{2}C_F^2\left(-\frac{3}{4} + \ln v - \ln\left(\frac{\mu_F^2}{s}\right)\right)\right. \\ &+ 3C_A^2\left(\ln\left(\frac{v}{1-v}\right) - \ln\left(\frac{\mu_F^2}{s}\right)\right) \\ &\left. + \frac{3}{2}C_F C_A\left(-\ln(1-v) + 3\ln v - 3\ln\left(\frac{\mu_F^2}{s}\right)\right.\right. \\ &\left.\left. - \frac{3}{2}\right) - \frac{\beta_0}{2}\left(\frac{C_F}{4} + C_A\right)\right]\left[\frac{\ln^2(1-w)}{1-w}\right]_+ \\ &\left. + \mathcal{O}\left(\left[\frac{\ln(1-w)}{1-w}\right]_+\right)\right\}\,, \end{aligned} \tag{8}$$

with $\beta_0 = (11C_A - 2n_f)/3$. We have also derived all NNLL terms in the NNLO expansion by matching with the exact NLO cross section [8]. Analogous results have been obtained for the partonic channel $q\bar{q} \to g\gamma$ [1, 8].



Figure 1. Exact NLO and approximate NNLO results for direct photon production.

In Fig. 1 we show some numerical results [8] for the direct photon production cross section as a function of the photon p_T and compare with the experimental results from the E706 Collaboration at Fermilab [14]. We see that the sum of the exact NLO cross section and NNLO approximate corrections shows a much reduced dependence on the factorization scale relative to the exact NLO cross section alone. However, the NNLO cross section is still below the E706 data.

3. W + jet production

In this section, we discuss resummation for W + jet production [1, 15] in hadronic processes

$$h_1 + h_2 \to W + X\,. \tag{9}$$

The construction of the resummed cross section is completely analogous to what we presented in the previous section for direct photon production, Eq. (4). The soft anomalous dimensions are the same

as for direct photon production. In this section we expand the NLL resummed cross section for W + jet production up to NNLO and perform a comparison with the NLO fixed-order results of Refs. [16, 17]. We give explicit results for the partonic subprocess

$$q(p_1) + \bar{q}(p_2) \to W(Q) + g(p_J)\,. \qquad (10)$$

The threshold region is given in terms of the variable $s_2 = s + t + u - Q^2$ by $s_2 = 0$, and we find plus distributions of the form $[\ln^m(s_2/Q^2)/s_2]_+$ at every order in perturbative QCD.

The NLO $\overline{\rm MS}$ soft gluon corrections for this partonic process at NLL accuracy are [1, 15]

$$\begin{aligned} E_Q \frac{d\hat{\sigma}^{\rm NLO}_{q\bar{q}\to gW}}{d^3Q} &= \sigma^B_{q\bar{q}\to gW} \frac{\alpha_s(\mu_R^2)}{\pi} \\ &\times \left\{ (4C_F - C_A) \left[\frac{\ln(s_2/Q^2)}{s_2}\right]_+ \right. \\ &+ \left[(2C_F - C_A) \ln\left(\frac{sQ^2}{tu}\right) - \frac{\beta_0}{4} \right. \\ &\left. \left. - 2C_F \ln\left(\frac{\mu_F^2}{Q^2}\right) \right] \left[\frac{1}{s_2}\right]_+ \right. \\ &\left. + \mathcal{O}(\delta(s_2)) \right\}\,. \qquad (11) \end{aligned}$$

Here the Born term is

$$\begin{aligned} \sigma^B_{q\bar{q}\to gW} &= \frac{\alpha\alpha_s(\mu_R^2) C_F}{sN_c} (|L_{f_2 f_1}|^2 + |R_{f_2 f_1}|^2) \\ &\times \left(\frac{u}{t} + \frac{t}{u} + \frac{2Q^2 s}{tu}\right)\,, \qquad (12) \end{aligned}$$

with L and R the left- and right-handed couplings of the W boson to the quark line of flavor f [17]. The NLO expansion is in agreement with the exact one-loop results in Ref. [17]. All $\delta(s_2)$ terms can be obtained by matching to the exact NLO result.

The NNLO $\overline{\rm MS}$ soft gluon corrections for the partonic process $q\bar{q} \longrightarrow gW$ are [15]

$$\begin{aligned} E_Q \frac{d\hat{\sigma}^{\rm NNLO}_{q\bar{q}\to gW}}{d^3Q} &= \sigma^B_{q\bar{q}\to gW} \left(\frac{\alpha_s(\mu_R^2)}{\pi}\right)^2 \\ &\times \left\{ \frac{1}{2}(4C_F - C_A)^2 \left[\frac{\ln^3(s_2/Q^2)}{s_2}\right]_+ \right. \\ &+ \left[-\frac{3}{2}(4C_F - C_A) \left(2C_F \ln\left(\frac{\mu_F^2}{Q^2}\right) \right. \right. \\ &\left. + (2C_F - C_A) \ln\left(\frac{tu}{sQ^2}\right) \right) \\ &\left. - \frac{\beta_0}{2}\left(5C_F - \frac{3}{2}C_A\right) \right] \left[\frac{\ln^2(s_2/Q^2)}{s_2}\right]_+ \\ &\left. + \mathcal{O}\left(\left[\frac{\ln(s_2/Q^2)}{s_2}\right]_+\right) \right\}\,. \qquad (13) \end{aligned}$$

We can also derive all NNLL terms in the NNLO expansion by matching with the exact NLO cross section [15]. Analogous results have been obtained for the partonic channel $qg \to qW$ [1, 15]. In future work we plan to study the numerical significance of resummation for $W+$ jet production.

This work was supported in part by the US Department of Energy.

References

[1] N. Kidonakis, invited review for *Int. J. Mod. Phys.* A, hep-ph/9902484.

[2] N. Kidonakis, in *JHEP Proceedings, Corfu Summer Institute on Elementary Paricle Physics, 1998*, hep-ph/9904507.

[3] N. Kidonakis, G. Oderda, and G. Sterman, in proceedings of the *EPIC'99 Workshop*, 1999, hep-ph/9906338.

[4] E. Laenen, G. Oderda, and G. Sterman, *Phys. Lett.* B **438** (1998) 173.

[5] S. Catani, M.L. Mangano and P. Nason, *JHEP* **9807** (1998) 024.

[6] S. Catani, M.L. Mangano, P. Nason, C. Oleari, and W. Vogelsang, *JHEP* **9903** (1999) 025.

[7] N. Kidonakis, in proceedings of the *DIS99 Workshop*, Zeuthen, April 19-23, 1999, hep-ph/9905480.

[8] N. Kidonakis and J. Owens, in preparation.

[9] G. Sterman, *Nucl. Phys.* **B281** (1987) 310.

[10] N. Kidonakis and G. Sterman, *Phys. Lett.* B **387** (1996) 867; *Nucl. Phys.* **B505** (1997) 321; N. Kidonakis, Ph.D. Thesis, hep-ph/9606474.

[11] N. Kidonakis, G. Oderda, and G. Sterman, *Nucl. Phys.* **B525** (1998) 299; **B531** (1998) 365.

[12] N. Kidonakis, J. Smith, and R. Vogt, *Phys. Rev.* D **56** (1997) 1553; N. Kidonakis, in *QCD 97*, *Nucl. Phys.* **B** *(Proc. Suppl.)* **64** (1998) 402; N. Kidonakis and R. Vogt, *Phys. Rev.* D **59** (1999) 074014.

[13] L.E. Gordon and W. Vogelsang, *Phys. Rev.* D **48** (1993) 3136.

[14] E706 Collaboration, L. Apanasevich *et al.*, *Phys. Rev. Lett.* **81** (1998) 2642.

[15] N. Kidonakis and V. Del Duca, in preparation.

[16] P.B. Arnold and M.H. Reno, *Nucl. Phys.* **B319** (1989) 37; (E) **B330** (1990) 284; *Phys. Rev.* D **40** (1989) 912.

[17] R.J. Gonsalves, J. Pawlowski and C.-F. Wai, *Phys. Rev.* D **40** (1989) 2245.